

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/275036925>

Forgotten records of *Chrysopelea taprobanica* Smith, 1943 (Squamata: Colubridae) from India

Article in *Zootaxa* · April 2015

DOI: 10.11646/zootaxa.3947.4.11

CITATIONS

0

READS

117

3 authors:



[Ruchira Somaweera](#)

CSIRO Land and Water

52 PUBLICATIONS 397 CITATIONS

[SEE PROFILE](#)



[Davi Lima Pantoja](#)

Universidade Federal do Piauí

15 PUBLICATIONS 31 CITATIONS

[SEE PROFILE](#)



[V. Deepak](#)

Indian Institute of Science

31 PUBLICATIONS 44 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Ecological and Behavioral traits of *Varanus salvator salvator* (Reptilia: Varanidae) in an suburban landscape of southwestern Sri Lanka [View project](#)



Systematics, diversification and biogeography of fan-throated lizards [View project](#)

All content following this page was uploaded by [V. Deepak](#) on 18 April 2015.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.



<http://dx.doi.org/10.11646/zootaxa.3947.4.11>

<http://zoobank.org/urn:lsid:zoobank.org/pub:B1A7C3D5-EF2D-4244-8DEC-C2B0DC29BB7F>

Forgotten records of *Chrysopelea taprobanica* Smith, 1943 (Squamata: Colubridae) from India

RUCHIRA SOMAWEERA¹, DAVI LIMA PANTOJA² & V. DEEPAK³

¹CSIRO Land and Water Flagship, 50B Angove St., North Perth, WA 6006. Australia

²Departamento de Zoologia, IB. Universidade de Brasília. Brasília, DF. 70910-900. Brazil

³TE-11, Center for Ecological Sciences, Indian Institute of Science, Bangalore – 560 012, Karnataka, India

⁴Corresponding author. E-mail: deepaksalea@gmail.com

The colubrid snake *Chrysopelea taprobanica* Smith, 1943 was described from a holotype from Kanthali (= Kantalai) and paratypes from Kurunegala, both localities in Sri Lanka (formerly Ceylon) (Smith 1943). Since its description, literature pertaining to Sri Lankan snake fauna considered this taxon to be endemic to the island (Taylor 1950, Deraniyagala 1955, de Silva 1980, de Silva 1990, Somaweera 2004, Somaweera 2006, de Silva 2009, Pyron *et al.* 2013). In addition, earlier efforts on the Indian peninsula (e.g. Das 1994, 1997, Das 2003, Whitaker & Captain 2004, Aengals *et al.* 2012) and global data compilations (e.g. Wallach *et al.* 2014, Uetz & Hošek 2015) did not identify any record from mainland India until Guptha *et al.* (2015) recorded a specimen (voucher BLT 076 housed at Bio-Lab of Seshachalam Hills, Tirupathi, India) in the dry deciduous forest of Chamala, Seshachalam Biosphere Reserve in Andhra Pradesh, India in November 2013. Guptha *et al.* (2015) further mentioned an individual previously photographed in 2000 at Rishi Valley, Andhra Pradesh, but with no voucher specimen collected. Guptha's record, assumed to be the first confirmed record of *C. taprobanica* in India, is noteworthy as it results in a large range extension, from northern Sri Lanka to eastern India with an Euclidean distance of over 400 km, as well as a change of status, *i.e.*, species not endemic to Sri Lanka. However, at least three little-known previous records of this species from India evaded most literature and were overlooked by the researchers including ourselves.

In 1949, John D. Constable mentioned specimens of *C. taprobanica* from Madras in India (Constable 1949). Constable (1949) was subsequently cited by Mertens (1968) in a paper published in German, and listed 'India' under the distribution of the taxon. Possibly referring to Constable's record, Mahendra (1984) listed 'Indian mainland near Madras' under the distribution of *C. taprobanica* along with Ceylon.

Constable (1949) reported a single voucher (number MCZ 47881: Fig. 1A–B); the collector name R. H. Beddome, and the scientific collection: Museum of Comparative Zoology (MCZ), Harvard University, USA. There is no date of collection associated with, but based on the museum's collection books, the voucher was first catalogued in 1876 (see the MCZ 47881 remarks: 'reregistered from MCZ 3903', which was catalogued in 1876). The specimen is a male with 17 midbody scale-rows, 208 ventrals, 120 subcaudals, divided anal scale, 9 upper labials, 1 loreal, 1 preocular, 2 postoculars, 2+2 temporals, and total length of 785 (560 + 225) mm (Constable 1949). Although Constable (1949) only mentioned one voucher number, he indicated that more than one specimen is associated with that number (e.g. '*CT. Head*' and '*these examples show that this species, originally described as from Ceylon alone, also occurs on the Indian mainland*'). In fact, a fore body of another *C. taprobanica* collected by Col. Richard H. Beddome presumably along with MCZ 47881, has been now recatalogued (in '18/xii/1992') as specimen MCZ 177292 (Fig. 1C–D) at the Museum of Comparative Zoology.

Noteworthy, there are two additional specimens associated to the aforementioned voucher MCZ 3903. The first one retains this voucher number and is indeed a *Chrysopelea ornata* (Shaw, 1802) (confirmed by us through photographs of the specimen), the other specimen was sent to the Coleção Herpetológica do Butantan, São Paulo, Brazil (see the MCZ 3903 remarks: 'To Butantan 5/31/29'). The latter specimen received the voucher IBSP-4273 and was catalogued as *C. ornata*. However, the specimen has not been recovered after the fire that destroyed most of the Butantan collection in 2010 (Giuseppe Puerto pers. comm.), therefore the identity has not been verified.

A third specimen of *C. taprobanica* with only 'India' as the collection locality is deposited at the Naturhistorisches Museum Wien (NMW: Natural History Museum of Vienna) under the voucher number NMW 27287:34 (G. Vogel pers. comm.: Fig. 1E–F). No date of collection exists for this specimen. This male specimen (snout vent length 395 mm)




FIGURE 1. Specimens of *Chrysopelta taprobanica* from India. A-B: MCZ 47881 and C-D: MCZ 177292 housed at the Museum of Comparative Zoology, Harvard; E-F: NMW 27287:34 housed at Natural History Museum, Vienna [photographed by Joseph Martinez (A-D) and Gernot Vogel (E-F)]. Scale bars = 1 cm. Pictures on the fourth known specimen from India (BLT 076) are available in Guptha *et al.* (2015).

currently labeled as *C. ornata* is indeed a *C. taprobanica* based on color pattern and scalation. The specimen has 17 mid-body dorsal scales, 204 ventrals (with preventral) and an incomplete tail (G. Vogel pers. comm.).

Despite differences in scalation, *C. taprobanica* was considered only a colour variation of *C. ornata* before Smith's description in 1943 (see Hora & Jayaram 1949). We have verified the identity of 13 such specimens of *C. ornata* collected prior to 1943 from India (Appendix A). However, it is possible that additional specimens of *C. taprobanica* collected prior to 1943 are lodged in other museums and evaded our notice. Because our compiled information from literature and scientific collections refer only to very wide regions (*i.e.*, India, or Madras), it is not possible to define a precise record locality, and therefore, the map of *C. taprobanica* records available at Guptha *et al.* (2015) remains as the most up-to-date of its known distribution of records.

The Madras region where the MCZ specimens were collected and the Seshachalam hill range where Guptha *et al.* (2015) recently recorded *C. taprobanica* is part of the Eastern Ghats. Eastern Ghats was generally considered as a poor sister of Western Ghats in terms of explorations and study (Das 2002, Srinivasulu & Das 2008). Recent explorations in the Eastern Ghats led to rediscovery of endemic species and discovery of new species (e.g. Agarwal *et al.* 2012, Aengals & Ganesh 2013, Agarwal *et al.* 2013, Datta-Roy *et al.* 2013). The lag of detailed explorations in their likely range in Southern part of Eastern Ghats till recently could also be a probable cause for the lack of additional sightings of *C. taprobanica* from India.

Acknowledgements

We thank Gernot Vogel for sharing his literature and museum records. Jonathan Losos, Joseph Martinez, Alex Pyron, Max A. Nickerson, Addison Wynn, Jens Vindum, Lauren Vonnahme, Christopher Raxworthy, Giuseppe Puerto, Noel Graham, Daniel Klingberg Johansson, Jon Fjeldså, and Pedro H. Bernardo assisted with obtaining photographs and details of museum specimens. Older literature was accessed through the Biodiversity Heritage Library.

References

- Aengals, R. & Ganesh, S.R. (2013) *Rhinophis goweri*- a new species of Shield-tailed snake from the Southern Eastern Ghats, India. *Russian Journal of Herpetology*, 20, 61–65.
- Aengals, R., Sathish Kumar, V.M. & Palot, M.J. (2012) Updated Checklist of Indian Reptiles. Available from: <http://zsi.gov.in/checklist/Checklist%20of%20Indian%20Reptiles.pdf> (accessed 31 March 2015)
- Agarwal, I., Dutta-Roy, A., Bauer, A. & Giri, V. (2012) Rediscovery of *Geckoella jeyporensis* (Squamata: Gekkonidae), with notes on morphology, coloration and habitat. *Hamadryad*, 36, 17–24.
- Agarwal, I., Wilkinson, M., Mohapatra, P.P., Dutta, S.K., Giri, V.B. & Gower, D.J. (2013) The first teresomatan caecilian (Amphibia: Gymnophiona) from the Eastern Ghats of India—a new species of *Gegeneophis* Peters, 1880. *Zootaxa*, 3693 (4), 534–546.
<http://dx.doi.org/10.11646/zootaxa.3693.4.7>
- Constable, J.D. (1949) Reptiles from India Peninsula in the museum of comparative Zoology. *Bulletin Museum of comparative Zoology*, 103, 59–160.
- Das, I. (1994) The reptiles of South Asia: checklist and distributional summary. *Hamadryad*, 19, 15–40.
- Das, I. (1997) Checklist of the reptiles of India, with English common names. *Hamadryad*, 22, 32–45.
- Das, I. (2002) *A photographic guide to snakes and other reptiles of India*. New Holland, Ralph Curtis books, Sanibel Island, 144 pp.
- Das, I. (2003) Growth of knowledge on the reptiles of India, with an introduction to systematics, taxonomy and nomenclature. *Journal of Bombay Natural History Society*, 100, 3.
- Datta-Roy, A., Mohapatra, P.P., Dutta, S.K., Giri, V.B., Veerappan, D., Maddock, S.T., Raj, P., Agarwal, I. & Karanth, P. (2013) A long-lost relic from the Eastern Ghats: Morphology, distribution and habitat of *Sepsophis punctatus* Beddome, 1870 (Squamata: Scincidae). *Zootaxa*, 3670, 055–062.
- de Silva, A. (1990) *Colour guide to the snakes of Sri Lanka*. R & A Publishing Ltd, Portishead, UK, 130 pp.
- de Silva, A. (2009) *Snakes of Sri Lanka: a coloured atlas (with accounts on snakes in archaeology, history, folklore, venomous snakes and snakebite)*. Vijitha Yapa Publications, Kohuwala, 135 pp.
- de Silva, P.H.D.H. (1980) *Snake Fauna of Sri Lanka: With Special Reference to Skull, Dentition, and Venom in Snakes*. National Museums of Sri Lanka, Colombo, 472 pp.
- Deraniyagala, P. (1955) *A Colored Atlas of some Vertebrates from Ceylon. Vol. III. Serpentine Reptilia*. The National

- Museum, Government Press, Colombo, Sri Lanka, xix + 121 pp.
- Guptha, B., Prasad, N.V.S., Maddock, S.T. & Deepak, V. (2015) First record of *Chrysopelea taprobanica* Smith, 1943 (Squamata: Colubridae) from India. *Check List*, 11, 1523.
<http://dx.doi.org/10.15560/11.1.1523>
- Hora, S.L. & Jayaram, K. (1949) Remarks on the distribution of snakes of Peninsular India with Malayan affinities. *Proceedings of the National Institute of Science*, 15, 399–402.
- Mahendra, B.C. (1984) *Handbook of the snakes of India, Ceylon, Burma, Bangladesh, and Pakistan*. Academy of Zoology, Agra, xvi, 412 pp.
- Pyron, R.A., Kandambi, H.K.D., Hendry, C.R., Pushpamal, V., Burbrink, F.T. & Somaweera, R. (2013) Genus-level phylogeny of snakes reveals the origins of species richness in Sri Lanka. *Molecular Phylogenetics and Evolution*, 66, 969–978.
<http://dx.doi.org/10.1016/j.ympev.2012.12.004>
- Smith, M.A. (1943) The Fauna of British India, Ceylon and Burma, Including the Whole of the Indo-Chinese Sub-Region. Reptilia and Amphibia. 3 (Serpentes). Taylor and Francis, London, 583 pp.
- Somaweera, R. (2004) Sri Lankan Colubrid Snakes. *Sri Lanka Naturalist*, 6, 32–46.
- Somaweera, R. (2006) Sri Lankawe Sarpayin ('The Snakes of Sri Lanka'). Wildlife Heritage Trust of Sri Lanka (WHT), 297 pp.
- Srinivasulu, C. & Das, I. (2008). The Herpetofauna of Nallamala Hills, Eastern Ghats, India: An Annotated Checklist, With Remarks on Nomenclature, Taxonomy, Habitat Use, Adaptive Types and Biogeography. *Asiatic Herpetological Research*, 11, 110–131.
- Taylor, E.H. (1950) A brief review of Ceylonese snakes. *University of Kansas Science Bulletin*, 33(14), 519–603.
- Uetz, P. & Hošek, J. (2015) The Reptile Database. Available from: <http://reptile-database.reptarium.cz/> (accessed 11 January 2015)
- Wallach, V., Williams, K.L. & Boundy, J. (2014) *Snakes of the World: A Catalogue of Living and Extinct Species*. CRC Press, Boca Raton, 1209 pp.
- Whitaker, R. & Captain, A. (2004) *Snakes of India*. Draco Books, Chennai. 479 pp.

APPENDIX A. Specimens of *Chrysopelea ornata* collected prior to 1943 from India whose identity was verified during the current study

American Museum of Natural History (New York, USA)—AMNH 43379, AMNH 43380, AMNH 74847; California Academy of Sciences (California, USA)—CAS 12388, CAS 17256, CAS 17257; Museum of Comparative Zoology (Cambridge, USA)—MCZ 3115, MCZ R3903; Royal Ontario Museum (Toronto, Canada)—ROM 16256; National Museum of Natural History (Washington, DC, USA)—USNM 129704; Natural History Museum of Denmark (Copenhagen, Denmark)—ZMUC R63722, ZMUC R63723, ZMUC R63738.